

**BASIC CARE DAYTIME NIGHTTIME COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride, guaifenesin  
L. Perrigo Company**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Amazon Daytime Nighttime Cold & Flu Drug Facts**

**Active ingredients (in each 30 mL) – Night Time**

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Doxylamine succinate 12.5 mg

Phenylephrine HCl 10 mg

**Purpose**

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

**Uses**

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- minor aches and pains
- headache
- fever
- sore throat
- runny nose and sneezing
- cough due to minor throat and bronchial irritation
- cough to help you sleep
- reduces swelling of nasal passages
- promotes nasal and/or sinus drainage
- temporarily restores freer breathing through the nose

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- sodium-restricted diet

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

**When using this product**

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

**Other information**

- **each 30 mL contains:** sodium 42 mg
- store at 20-25°C (68-77°F)

**Inactive ingredients**

alcohol, anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C blue #1, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose

**Questions or comments?**

**1-800-719-9260**

**Active ingredients (in each 30 mL) - Day Time**

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Guaifenesin 400 mg

Phenylephrine HCl 10 mg

## **Purpose**

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

## **Uses**

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- cough due to minor throat and bronchial irritation
- minor aches and pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

## **Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

## **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after

stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

**When using this product**

**do not use more than directed**

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
---------------------------------	-------------------

children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

## Other information

- **each 30 mL contains:** sodium 41 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

## Inactive ingredients

alcohol, anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C blue #1, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose

## Questions or comments?

**1-800-719-9260**

## Package/Label Principal Display Panel

Combo Pack!

Severe

Compare to Vicks® DayQuil® Severe+ VapoCOOL™ active ingredients

Non-Drowsy

daytime

cold & flu

Acetaminophen, Phenylephrine HCl, Dextromethorphan HBr, Guaifenesin

Pain Reliever/Fever Reducer

Cough Suppressant, Expectorant, Nasal Decongestant

ALCOHOL 10%

Vapor Ice™

12 FL OZ (355 mL)

Severe

Compare to Vicks® NyQuil® Severe+ VapoCOOL™ active ingredients

nighttime

cold & flu

Acetaminophen, Phenylephrine HCl, Doxylamine Succinate, Dextromethorphan HBr

Pain Reliever, Fever Reducer, Cough Suppressant, Antihistamine, Nasal Decongestant

ALCOHOL 10%

Vapor Ice™

12 FL OZ (355 mL)

NDC 0113-7280-02

basic+care®

**Combo Pack!**

**PARENTS:**  
Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

**DO NOT USE IF PRINTED MEDICATION IS  
BROKEN OR MISSING**  
\*These products are not manufactured or  
distributed by Procter & Gamble, distributor  
of Vicks® DayQuil® Severe+ VapoCOOL™  
and Vicks® NyQuil® Severe+ VapoCOOL™.

**Drug Facts (continued)**  
**Inactive ingredients:** alcohol, anhydrous citric acid, D&C yellow #10,  
edetate disodium, FD&C blue #1, flavor, glycerin, polyethylene glycol,  
propylene glycol, purified water, saccharin sodium, sodium benzoate,  
sodium chloride, sodium citrate, sorbitol solution, sucralose  
**Questions or comments?** 1-800-719-9260

**Severe**

Compare to  
Vicks® DayQuil®  
Severe+ VapoCOOL™  
active ingredients\*

Non-Drowsy

**daytime  
cold & flu**

**Acetaminophen**, Phenylephrine HCl,  
Dextromethorphan HBr, Guaifenesin

Pain Reliever/Fever Reducer,  
Cough Suppressant, Expectorant,  
Nasal Decongestant

ALCOHOL 10%



Vapor Ice™



12 FL OZ (355 mL)

**Severe**

Compare to  
Vicks® NyQuil®  
Severe+ VapoCOOL™  
active ingredients\*

**nighttime  
cold & flu**

**Acetaminophen**, Phenylephrine HCl,  
Doxylamine Succinate, Dextromethorphan HBr

Pain Reliever, Fever Reducer,  
Cough Suppressant, Antihistamine,  
Nasal Decongestant

ALCOHOL 10%



Vapor Ice™



12 FL OZ (355 mL)

Distributed by:  
Perrigo  
Allergan, MI 49010

**GLUTEN FREE**

**Drug Facts (continued)**  
**Inactive ingredients:** alcohol, anhydrous citric acid, D&C yellow #10,  
edetate disodium, FD&C blue #1, flavor, glycerin, polyethylene glycol,  
propylene glycol, purified water, saccharin sodium, sodium benzoate,  
sodium chloride, sodium citrate, sorbitol solution, sucralose  
**Questions or comments?** 1-800-719-9260

Drug Facts	Nighttime	Drug Facts	Daytime
<b>Active ingredients (in each 30 mL)</b> Acetaminophen 650 mg.....Pain reliever/fever reducer Dextromethorphan HBr 20 mg.....Cough suppressant Doxylamine succinate 12.5 mg.....Antihistamine Phenylephrine HCl 10 mg.....Nasal decongestant		<b>Active ingredients (in each 30 mL)</b> Acetaminophen 650 mg.....Pain reliever/fever reducer Dextromethorphan HBr 20 mg.....Cough suppressant Guaifenesin 400 mg.....Expectorant Phenylephrine HCl 10 mg.....Nasal decongestant	
<b>Uses</b> temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ sinus congestion and pressure ■ minor aches and pains ■ headache ■ sore throat ■ runny nose and sneezing ■ cough to help you sleep ■ cough due to minor throat and bronchial irritation ■ reduces swelling of nasal passages ■ fever ■ promotes nasal and/or sinus drainage ■ temporarily restores freer breathing through the nose		<b>Uses</b> temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ sinus congestion and pressure ■ cough due to minor throat and bronchial irritation ■ minor aches and pains ■ headache ■ fever ■ reduces swelling of nasal passages ■ temporarily restores freer breathing through the nose ■ sore throat ■ helps to open phlegm (mucus) and thin bronchial secretions to rid the bronchial passages away of bothersome mucus and make coughs more productive ■ promotes nasal and/or sinus drainage	
<b>Warnings</b> <b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product <b>Allergy alert:</b> Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away. <b>Sore throat warning:</b> If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. <b>Do not use</b> ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients		<b>Warnings</b> <b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product <b>Allergy alert:</b> Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away. <b>Sore throat warning:</b> If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. <b>Do not use</b> ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients	
<b>Ask a doctor before use if you have</b> ■ liver disease ■ heart disease ■ high blood pressure ■ a breathing problem such as emphysema or chronic bronchitis ■ thyroid disease ■ diabetes ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ a sodium-restricted diet		<b>Ask a doctor before use if you have</b> ■ liver disease ■ heart disease ■ thyroid disease ■ high blood pressure ■ trouble urinating due to an enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ diabetes ■ a sodium-restricted diet	
<b>Ask a doctor or pharmacist before use if you are</b> ■ taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin		<b>Ask a doctor or pharmacist before use if you are</b> ■ taking the blood thinning drug warfarin	
<b>When using this product</b> ■ do not use more than directed ■ excitability may occur, especially in children ■ be careful when driving a motor vehicle or operating machinery ■ marked drowsiness may occur ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ avoid alcoholic drinks		<b>When using this product</b> do not use more than directed	
<b>Stop use and ask a doctor if</b> ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ you get nervous, dizzy or sleepless ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.		<b>Stop use and ask a doctor if</b> ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.	
<b>If pregnant or breast-feeding</b> , ask a health professional before use. <b>Keep out of reach of children.</b> <b>Overdose warning:</b> In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.		<b>If pregnant or breast-feeding</b> , ask a health professional before use. <b>Keep out of reach of children.</b> <b>Overdose warning:</b> In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.	
<b>Directions</b> ■ take only as directed—see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs adults & children 12 yrs & over 30 mL every 4 hrs children 4 to under 12 yrs ask a doctor children under 4 yrs do not use		<b>Directions</b> ■ take only as directed—see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs adults & children 12 yrs & over 30 mL every 4 hrs children 4 to under 12 yrs ask a doctor children under 4 yrs do not use	
<b>Other information</b> ■ each 30 mL contains: sodium 42 mg ■ store at 20-25°C (68-77°F)		<b>Other information</b> ■ each 30 mL contains: sodium 41 mg ■ store at 20-25°C (68-77°F). Do not refrigerate.	

10 284092 :



## BASIC CARE DAYTIME NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride, guaifenesin kit

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0113-7280
--------------	----------------	--------------------	---------------

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0113-7280-02	1 in 1 PACKAGE; Type 0: Not a Combination Product	02/25/2020	

### Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE	355 mL
Part 2	1 BOTTLE	355 mL

### Part 1 of 2



# BASIC CARE NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride solution

## Product Information

Route of Administration ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9B19B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

## Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341		

Part 2 of 2

BASIC CARE DAYTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution

Product Information

Item Code (Source)	NDC:0113-7902
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 30 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0113-7902-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	02/25/2020	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	02/25/2020	

**Labeler** - L. Perrigo Company (006013346)

Revised: 4/2020

L. Perrigo Company